

Religiosity and health-related quality of life in HIV positive young adults seeking health care in Alebtong District, Northern Uganda: A cross-sectional study

Sarah Nabuwufu¹, Augustine Bilitya², Binepe Bwambale Lhwanzo³, Mariam Nakazibwe⁴, Gloria Aleso⁵, Fazira Namata⁶, and Amir Kabunga⁷

Abstract:

Objectives: This study estimated religiosity and health-related quality of life in HIV positive young adults aged 18 to 36 years seeking care in Alebtong District Northern Uganda.

Methods: A descriptive, cross-sectional design was used among 363 HIV-positive young adults in five health centers in Alebtong district, northern Uganda in April 2021.

Results: Positive religiosity is associated with better physical health ($\beta= 0.52$, $P= 0.01$) and psychological health ($\beta= 0.63$, $P= 0.02$). Negative religiosity was associated with poor Health Related Quality of Life (HRQOL) domains of physical health ($\beta= -0.43$, $P= 0.02$), psychological health ($\beta= -0.75$, $P= 0.02$), social relationship health ($\beta= -0.53$, $P= 0.04$) and environmental health ($\beta=-0.66$, $P= <0.01$)

Conclusion: The findings in this study show that religiosity is significantly associated with HRQOL. This underscores the importance of religiosity in the management of HIV-positive patients. Religiosity should be an integral part of the HIV treatment program to the HRQOL of HIV patients. The findings also suggest that HIV-positive patients are likely to benefit from support in the community. Thus, there is a need to strengthen the collaboration between religious organizations in the management of HIV to improve the HRQOL.

Keywords: *HIV, health facilities, health quality of life, religiosity, young adults*

INTRODUCTION

Global trends in HIV infection show that HIV remains a major global public health concern and has claimed 33 million lives by 2019 (Ng et al., 2020). Sub-Saharan Africa has a disproportionate liability of HIV, accounting for 71% of the global burden of infection (Ng et al., 2020). Out of an estimated daily 6000 new infections occurring globally, two out of three is in Sub-Saharan Africa with young adults bearing a disproportionate burden. (Kimera et al., 2020). HIV prevalence is nearly three times higher in young adults compared to those aged 15 to 19 (UNAIDS, 2019). While between 2000 and 2019, new HIV infections declined by 39% and HIV-related deaths reduced by 51%, it continues to be a major cause of morbidity affecting the health-related quality of life (HRQOL) of persons living with HIV (Engelhard et al., 2018). To improve the HRQOL, the World Health Organization (WHO) emphasizes that religiosity be an integral part of the treatment and management of HIV (Group, 1995). However, although several studies have examined the HRQOL in patients with HIV, only a few have examined the influence of religiosity on HRQOL among HIV-positive young adults especially in developing countries (Mutabazi-Mwesigire, Katamba, Martin, Seeley, & Wu, 2015).

¹²³⁴⁵⁶⁷ Faculty of Health Sciences, Lira University, P.O BOX 1035, Lira city

Correspondence may be sent to Amir Kabunga via email at amirkabs2017@gmail.com

26 Religiosity and health-related quality of life in HIV positive young adults

Religiosity is a person's belief, spirituality, and relevance toward a divinity (Gallagher & Tierney, 2013). In this study, religiosity was operationally defined as meaning, peace, and faith (Peterman, Fitchett, Brady, Hernandez, & Cella, 2002) among HIV-positive young adults. Available literature especially in developed countries indicates that individuals who have positive religiosity tend to report higher CD4 counts, low sex risk behaviours, decreased viral loads, and greater odds of HIV testing (Carrico et al., 2017). HIV patients use religiosity as a source of hope and comfort to cope with the disease and the weaknesses it causes. (Ferreira, Favoreto, & Guimarães, 2012) On the contrary, negative religiosity can affect the development of the disease and HRQOL (Hasanah¹, Ibrahim, & Sriati, 2019). The study by Szaflarsk and colleagues showed that one-third of the HIV patients' HRQOL in the sample reported improved due to positive religiosity among other factors (Szaflarski et al., 2006). Mrus and colleagues (Mrus et al., 2006) suggested that religiosity could be a potentially fruitful target for intervention to improve HRQOL. However, some studies indicate that negative religiosity could have negative aspects on HRQOL (Folkman, 2008). Some HIV patients appeared to rely more on religiosity than biomedical treatment (Folkman, 2008), but, how widely spread this issue could be among HIV-positive patients remains unknown and how it could impact HRQOL, especially in low-income countries. Therefore, there is a need to understand the influence of religiosity on HRQOL among HIV young adults, especially in developing countries.

HRQOL refers to how an individual functions in their life, and their perceived wellbeing in mental, physical, and social aspects of health (Killewo, Heggenhougen, & Quah, 2010). According to the World Health Organization (WHO), HRQOL is a collection of different elements concerning an individual and their environment including how they perceive their goal, expectations, and concerns (Group, 1995). A new "beyond viral suppression" model adds a "fourth 90" to the UNAID; the 90-90-90 target to ensure that 90% of people living with HIV with viral load suppression have a good HRQOL (Ahmed et al., 2021). Earlier studies have indicated factors influencing HRQOL including socio-demographics, stigma, CD4 T lymphocytes count, poor ART adherence, mental health, social support, adherence to ART, psychological status, and religiosity (Ahmed et al., 2021; Anuradha, Makkar, Nandi, & Rajeshwari, 2020). However, these factors may vary from one setting to another and from one culture to another (Ahmed et al., 2021).

The 2016 to 2017 Uganda AIDS Indicator Survey reported HIV prevalence in the northern region, Alebtong district inclusive to be at 7.2% which is higher than the national rate of 6.2% (Benyumiza et al., 2021). This may be attributed to the two decades of war (1986 to 2006) in the region involving widespread atrocities, which include abduction, forced marriages, mass rapes, and displacement (Kabunga, Anyolitoh, & Betty, 2020). Thus, although the war ended in 2006, it continues to impact the health and wellbeing of the post-war conflict-affected people (Katamba et al., 2020). Research has been done to assess the HRQOL in HIV patients and other people with chronic diseases concerning psychological health (Amoako-Agyeman, 2012). However, research on HRQOL of young adults living with HIV remains limited especially in developing countries. Besides, a review of Ugandan literature shows that very limited research has been conducted to assess the relationship between religiosity and HRQOL in HIV-positive young adults. To address this gap, this study investigated religiosity and HRQOL in HIV-positive young adults seeking care in Alebtong district health facilities in northern Uganda.

METHODOLOGY

Study design and settings

A cross-sectional study was conducted among HIV-positive young adults in five health centres in Alebtong district, northern Uganda in April 2021. Alebtong district is located in the Lango sub-region in northern Uganda. The district is bordered by Dokolo district to the south, Amuria district to the east, Otuoke district to the north, and Lira district to the west. This is approximately 387kms by road, north of Kampala, the capital city of Uganda. The HIV prevalence remains concentrated in the northern part of Alebtong, which may be due to the impact of the two decades of war (1986 to 2006) in the region involving widespread atrocities that include abduction, forced marriages, mass rapes, and displacement. Although the war ended in 2006, it continues to impact the health and wellbeing of the post-war conflict-affected people (Katamba et al., 2020).

Study Participants

The study population involved young adults aged 18 to 36 years attending Alebtong health centres. Statistics on HIV/AIDS in Uganda show that close to 1200 HIV-positive young adults lived in the Alebtong district in 2019 (Benyumiza et al., 2021). The sample size of the participants was calculated using the formula for finite population (Reid & Boore, 1991) where n= a sample size, N= the population size, and e = significance level, taking alpha as 0.05. Thus, the sample size was 300. However, more participants above the minimum sample size were included in the study. A consecutive sampling technique to select the representative sample was also employed in this study. Given the nature and availability of the respondents, every participant meeting the inclusion criteria was selected until the required sample size was achieved. However, patients who were severely ill were excluded from the study.

Patients and Public involvement

No patients or members of the public were involved in this study

Study Instruments

The researchers' generated questionnaire was used to collect data on socio-demographics of the participants including age, gender, marital status, religious affiliation and, level of education. Religiosity was assessed using the Duke University Religion Index (DUREL). The five-item scale assesses three major dimensions of Organizational Religious -Activity (ORA), Non-Organizational Religious Activity (NRA), and Intrinsic Religiosity (IR) (Young & Clark, 2013). Organizational and Non-Organizational items were rated on a six point-Likert scale. Intrinsic Religiosity consisted of three items on a five-point Likert scale. The sub-scales were examined indecently, with a higher score indicating higher ORA, NRA, and IR (Young & Clark, 2013). The DUREL has been validated in both clinical and community settings (Young & Clark, 2013).

The World Health Organization Quality of Life BREF (WHOQOL- BREF) was administered to collect data on HRQOL. The WHOQOL- BREF is a 26-item test rated on a five-point Likert scale, ranging from 1 (not at all to five (completely) (Meemon, Paek, Yenchai, & Wan, 2016). The instrument has been validated among people living with HIV in Uganda (Ndikuno, Namutebi, Kuteesa, Mukunya, & Olwit, 2016). The 24 facets of WHOQOL- BREF measure four domains of a person's perception within the last two weeks namely: psychological health, physical health, relationship with one's environment, and social relationships (Meemon et al., 2016). Two items assess the overall quality of life and satisfaction with general health. The domain scores were calculated and transformed using WHOQOL- BREF SPSS syntax file. We transformed scores on a scale ranging from 0 to 100 with higher scores indicating a better quality of life and vice-versa (Organization, 1996). In this study, the Cronbach alpha values were 0.88 for physical health, 0.75 for psychological health, 0.83 for environmental health, and 0.71 for social relationships.

Procedure

The study participants were young adults, ages 18 to 36 years old, attending Alebtong health centres. Five research assistants with data collection experience were recruited to collect data, and they were under the supervision of the principal investigator. A one-day training was given to data collectors to help them know the purpose of the study and the rights of the respondents to either decline or participate. Data collection was carried out for 10 days in April 2021. Each participant with HIV attending health centres and meeting the criteria was invited to participate in the study. Those who agreed to participate were informed about the purpose of the study. The informed consent document was read to the potential participant and signed after agreeing to participate.

Data analysis

Data were summarized by descriptive statistics including frequency, means, standard deviation, ranges, and percentages. The relationships between social-demographics factors and religiosity and HRQOL were assessed. An ANOVA analysis using the Bonferroni test was also performed to establish the relationship between independent groups. A multiple linear regression test was conducted to assess the correlation of components of religiosity and HRQOL after controlling for social-demographics factors. For all statistical tests, a significant level of P≤.05 was used. SPPSS for windows version 23 was also used to perform statistical analyses.

RESULTS

Results in Table 1 show that of the total 363 respondents, 53.7% were female, 74.6% were single, 46.5% had A-Level education, 34.7% were Catholics, 20.9% were Muslims, 29.7% were Protestants, 14.0% belonged to others and young (mean age, 33.3 ± 10.7), 26.9% of the participants reported that religion was related to treatment. Regarding religion and treatment, 26.7% believed it was indispensable.

Table 1: Socio-demographic characteristics of 363 HIV positive young adults

	Mean/Frequency	Std/Percentages
Age (years)	33.3	10.7
Gender		
Female	195	53.7
Male	168	46.2
Education level		
Primary education	10	2.7
O-Level	92	25.3
A-Level	169	46.5
Post-secondary	92	25.3
Marital status		
Single	271	74.6
Married	72	19.8
Widowed/Separated	20	5.5
Religious affiliation		
Catholics	126	34.7
Muslims	76	20.9
Protestants	108	29.7
Others	52	14.3
Religion and treatment		
Not needed	55	15.1
Somehow needed	59	16.2
Needed	77	21.2
Much needed	74	20.3
Indispensable	98	26.9

A-Level = Advanced Secondary Level of Education, O-Level= Ordinary Secondary Level of Education

Results in Table 2 show that religiosity was 17.5 ± 6.9 for positive religiosity and 11.4 ± 4.4 for negative religiosity. Respondents reported mean scores of 3.4 ± 1.3 for ORA, 4.1 ± 1.8 for NRA, and 6.6 ± 3.5 for IR. The transformed scores for the WHOL-BREF scale indicate that respondents reported 63.2 ± 15.5 for physical health, 55.1 ± 18.1 for psychological health, 57.9 ± 21.1 for social relationships, and 61.5 ± 16.6 , for environmental health. For overall HRQOL the respondents reported a mean score of 3.54 ± 0.8 .

Table 2: Religiosity, religious commitment, and HRQOL domains

Variable	Mean	SD
Religiosity		
Positive	17.5	6.8
Negative	11.4	4.4
Religious commitment		
ORA	3.4	1.3
NRA	4.1	1.8
IR	6.6	3.5
HRQOL		
Physical health	64.2	15.5
Psychological health	55.1	18.1
Social relationships health	57.9	21.1
Environment health	61.5	16.6
Overall HRQOL	3.54	0.8

A post-hoc correction analysis using the Bonferroni test (Table 3) shows that under DUREL, the highest mean scores were reported among the Catholics (4.3 ± 1.4) and IR (11.0 ± 2.8). The Muslims reported the highest positive religiosity (23.4 ± 5.4).

Table 3: Religiosity, religious commitment, and HRQOL domains

Mean (SD)	Religious affiliation			
	Catholics	Protestants	Islam	P-Value
Religious commitment				
ORA	4.3 ± 1.4	3.0 ± 1.1	3.7 ± 1.2	
NRA	3.5 ± 1.8	2.3 ± 1.4	3.6 ± 1.7	<0.01
IR	11.0 ± 2.8	9.5 ± 3.0	11.6 ± 3.2	
Religiosity				
Positive	20.4 ± 5.7	15.6 ± 6.2	23.4 ± 5.4	<0.01
Negative	12.9 ± 5.2	11.2 ± 4.7	14.1 ± 6.1	
HRQOL				
Physical	64.8 ± 15.8	65.0 ± 15.5	60.6 ± 18.6	
Psychological	57.0 ± 18.1	55.2 ± 18.1	56.7 ± 21.7	<0.01
Social relationships	57.5 ± 19.9	58.4 ± 17.3	60.3 ± 23.5	
Environment	63.5 ± 18.0	62.8 ± 13.8	59.2 ± 20.6	

Relationship between religiosity and HRQOL domains

A multiple linear regression model of positive and negative religiosity in HRQOL domains was performed (see Table 4). The results indicate that positive religiosity is associated with better physical health ($\beta= 0.52$, $P= 0.01$; 95%CI: 0.00-0.11) and psychological health ($\beta= 0.63$, $P= 0.02$; 95% CI: 1.42-3.33). Negative religiosity was associated with poor HRQOL domains of physical health ($\beta= -0.43$, $P= 0.02$; 95%CI: 4.07-11.11), psychological health ($\beta= -0.75$, $P= 0.02$; 95%CI: 3.16-7.82), social relationship health ($\beta= -0.53$, $P= 0.04$; 95%CI: 0.00-0.11) and environmental health ($\beta= -0.66$, $P= <0.01$; 95%CI: 0.00-0.11).

Table 4: Religiosity and HRQOL of HIV positive young adults (18-36 years)

Variables	Physical health				Psychological health				Social relationship health				Environmental health			
	B	SE	T	P	B	SE	T	P	B	SE	T	P	B	SE	T	P
Religiosity																
Positive	0.52	0.21	2.28	0.01	0.63	0.27	2.46	0.02	0.18	0.29	0.61	0.55	0.46	0.23	1.88	0.07
Negative	-0.43	0.19	-2.24	0.02	-0.75	0.22	-3.29	0.02	- 0.53	0.23	- 2.20	0.04	- 0.66	0.20	-3.12	<0.01
Religion in treatment																
Needed	Ref				Ref				Ref				Ref			
Not needed	1.98	2.49	0.80	0.44	7.08	2.92	2.79	0.01	4.65	3.09	1.53	0.14	4.37	2.63	1.65	0.11
Religious commitment																
ORA	1.05	0.76	1.34	0.17	2.46	0.93	2.71	0.01	2.65	0.96	2.75	0.01	2.10	0.81	2.54	0.01
NRA	-0.46	0.63	-0.72	0.47	-1.09	0.74	-1.45	0.16	- 1.32	0.79	- 1.66	0.11	- 0.44	0.67	-0.66	0.44
IR	-0.43	0.38	-0.06	0.26	-0.41	0.46	-0.91	0.35	0.84	0.48	1.74	0.07	0.12	0.41	0.33	0.65
Religious affiliation																
Catholics	Ref				Ref				Ref				Ref			
Muslims	-2.29	2.66	-0.84	0.38	-2.24	3.13	-0.06	0.92	5.07	3.32	1.52	0.22	- 3.02	2.84	-1.07	0.26
Protestants	3.04	2.44	1.24	0.23	2.10	2.90	1.01	1.29	5.19	3.04	1.72	0.08	3.42	2.61	1.31	0.20
Others	-0.97	2.69	-0.25	0.79	4.13	4.31	0.94	0.34	10.3	4.60	2.27	0.03	6.88	3.10	1.75	0,07

DISCUSSION

Religiosity and HRQOL in HIV Positive young adults aged 18-36 years seeking care at health facilities in Alebtong district, northern Uganda was estimated. Results of our study indicate that positive religiosity was related to better physical and psychological health. Conversely, negative religiosity was related to poor HRQOL domains of physical health, psychological health, social relationship health, and environmental health. The results suggest that religiosity influences the individual's wellbeing. 15.1% of HIV- positive HIV positive young adults attending Alebtong health centres reported that religiosity in treatment is not needed, it is evident from this study that HIV- positive patients believed that religiosity should be an integral part of HIV management. Additionally, our results suggest that negative religiosity may explain the poor HRQOL experienced by HIV-positive patients. This finding is important for HIV treatment programs because impaired HRQOL is believed to predict survival (de Boer-van der Kolk et al., 2010). Consistent with this view, Mrus and colleagues(Mrus et al., 2006) suggested that numerous correlates including religiosity could be potentially fruitful targets for intervention to improve HRQOL. It is therefore important to attend to negative religiosity to increase the odds of improving HRQOL in HIV patients.

A large body of research studies shows that positive religiosity has a positive role in people living with HIV including happiness and peace, fostering inner strength, acceptance, finding purpose in life, maintaining hope, and establishing relationships (Greene & Yoon, 2004). Although the small sample size in a few health centres reduces the generalizability of the results, the findings in our study are comparable to other studies (Ironson, Kremer, & Lucette, 2016). These studies indicate that patients with positive religiosity had higher CD4 cells and lower viral loads compared to patients whose levels of religiosity lessened post-diagnosis (Ironson et al., 2016). Our results mirror prior studies which report a positive relationship between religiosity and HRQOL (Desyani, Waluyo, & Yona, 2019). Other studies have also found similar results, revealing the correlation between religiosity and HRQOL (Abdala, Kimura, Koenig, Reinert, & Horton, 2015).

Our findings revealed that higher scores for ORA and IR were reported among the Christian participants. The Muslim participants in the current study reported higher positive and negative religiosity compared to other religious denominations. The findings in this study also indicate that involvement in religious activity was associated with better psychological and environmental domains of HRQOL. Our findings tend to suggest that religious community activities are beneficial to individuals living with HIV/AIDS. Thus, there is a need to strengthen the collaboration between religious organizations in the management of HIV to improve the HRQOL. While some faith-based organizations do not promote HIV-prevention or transmission activities like condom use, experience in some countries shows that in places where religious leaders have decided not to openly oppose them, there is a significant reduction in HIV infections and management.(Sidibé) Our results are in tandem with other studies assessing religiosity and HRQOL among HIV patients in Iran (Khademi et al., 2021).

CONCLUSION

The findings in this study show that religiosity is significantly associated with HRQOL. This underscores the importance of religiosity in the management of HIV-positive patients. Religiosity should be an integral part of the HIV treatment program to the HRQOL of HIV patients. The findings also suggest that HIV-positive patients are likely to benefit from support in the community. Thus, there is a need to strengthen the collaboration between religious organizations in the management of HIV to improve the HRQOL.

Acknowledgement: The authors wish to acknowledge the participants included in this study.

CONFLICT OF INTERESTS

The author(s) declared no potential conflict of interest concerning the research and/or publication of the article.

CONTRIBUTORS

SN, AB, BBL, MN, GA, LN planned the study, conducted the survey, performed the statistical analysis, and wrote the article. AK prepared the questionnaire, wrote the methodology, reviewed, edited, and supervised this work. SN helped in project administration and funding acquisition. All authors contributed to the final version of the article for publication.

FUNDING

'Research reported in this publication was supported by the Fogarty International Center (U.S. Department of State's Office of the U.S. Global AIDS Coordinator and Health Diplomacy [S/GAC] and the President's Emergency Plan for AIDS Relief [PEPFAR] of the National Institutes of Health under Award Number R25TW011210. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

ETHICAL CONSIDERATIONS

Ethical approval was obtained from the Institutional Review Board (IRB) of Gulu University (GURE-2021-38) and the Uganda National Council for Science and Technology. All participants provided written informed consent and received no facilitation for participation in this study. Participation was voluntary and no form of coercion was used. The data collected was confidential and anonymous with no information like names linking the study participants to the data. The COVID-19 protocols were strictly maintained throughout data collection. The respondents without masks were provided with masks. Sanitisers were available and social distancing was maintained.

REFERENCES

- Abdala, G. A., Kimura, M., Koenig, H. G., Reinert, K. G., & Horton, K. (2015). Religiosity and quality of life in older adults: literature review. *Life Style*, 2(2), 25-51.
- Ahmed, A., Saqlain, M., Bashir, N., Dujaili, J., Hashmi, F., Mazhar, F., . . . Awaisu, A. (2021). Health-related quality of life and its predictors among adults living with HIV/AIDS and receiving antiretroviral therapy in Pakistan. *Quality of Life Research*, 30(6), 1653-1664.
- Amoako-Agyeman, K. N. (2012). Adolescent religiosity and attitudes to HIV and AIDS in Ghana. *SAHARA-J: Journal of Social Aspects of HIV/AIDS*, 9(4), 227-241.
- Anuradha, S., Makkar, A. M., Nandi, P. K., & Rajeshwari, K. (2020). Quality of life and its determinants among people living with HIV attending an antiretroviral treatment center in Delhi, India. *Indian Journal of Public Health*, 64(2), 148.
- Benyumiza, D., Amongin, J. F., Ochaba, I., Adupa, M., Abuch, N., Banula, C. B., & Udho, S. (2021). Factors Associated with Utilization of HIV Testing Services among Adolescents Aged 10-19 Years in Lira District, Northern Uganda: A Cross-Sectional Study. *BioMed Research International*, 2021.
- Carrico, A. W., Storholm, E. D., Flentje, A., Arnold, E. A., Pollack, L. M., Neilands, T. B., . . . Johnson, W. (2017). Spirituality/religiosity, substance use, and HIV testing among young black men who have sex with men. *Drug and alcohol dependence*, 174, 106-112.
- de Boer-van der Kolk, I. M., Sprangers, M. A., Prins, J. M., Smit, C., de Wolf, F., & Nieuwkerk, P. T. (2010). Health-related quality of life and survival among HIV-infected patients receiving highly active antiretroviral therapy: a study of patients in the AIDS Therapy Evaluation in the Netherlands (ATHENA) Cohort. *Clinical infectious diseases*, 50(2), 255-263.
- Desyani, N. L. J., Waluyo, A., & Yona, S. (2019). The relationship between stigma, religiosity, and the quality of life of HIV-positive MSM in Medan, Indonesia. *Enfermería Clínica*, 29, 510-514.
- Engelhard, E. A., Smit, C., Van Dijk, P. R., Kuijper, T. M., Wermeling, P. R., Weel, A. E., . . . Nieuwkerk, P. T. (2018). Health-related quality of life of people with HIV: an assessment of patient related factors and comparison with other chronic diseases. *Aids*, 32(1), 103-112.
- Ferreira, D. C., Favoreto, C. A. O., & Guimarães, M. B. L. (2012). A influência da religiosidade no conviver com o HIV. *Interface-Comunicação, Saúde, Educação*, 16(41), 383-394.
- Folkman, S. (2008). The case for positive emotions in the stress process. *Anxiety, stress, and coping*, 21(1), 3-14.
- Gallagher, S., & Tierney, W. (2013). Religiousness/religiosity. *Encyclopedia of behavioral medicine*, 1653-1654.
- Greene, K. V., & Yoon, B. J. (2004). Religiosity, economics and life satisfaction. *Review of Social Economy*, 62(2), 245-261.
- Group, W. (1995). The World Health Organization quality of life assessment (WHOQOL): position paper from the World Health Organization. *Social science & medicine*, 41(10), 1403-1409.
- Hasanah¹, U., Ibrahim, K., & Sriati, A. (2019). Effects of Spiritual Counseling on Spiritual Health-Quality of Life in Patients with HIV/AIDS. *Nurse Media Journal of Nursing*, 13.

- Ironson, G., Kremer, H., & Lucette, A. (2016). Relationship between spiritual coping and survival in patients with HIV. *Journal of general internal medicine*, 31(9), 1068-1076.
- Kabunga, A., Anyolitho, M. K., & Betty, A. (2020). Emotional intelligence and compassion fatigue among psychotherapists in selected districts of Northern Uganda. *South African Journal of Psychology*, 50(3), 359-370.
- Katamba, A., Ogwang, M. D., Zamar, D. S., Muyinda, H., Oneka, A., Atim, S., . . . Friedman, A. J. (2020). Cango Lyec (healing the elephant): HIV incidence in post-conflict Northern Uganda. *EClinicalMedicine*, 23, 100408.
- Khademi, N., Zanganeh, A., Saeidi, S., Teimouri, R., Khezeli, M., Jamshidi, B., . . . Kiaee, K. G. (2021). Quality of life of HIV-infected individuals: insights from a study of patients in Kermanshah, Iran. *BMC Infectious Diseases*, 21(1), 1-12.
- Killewo, J., Heggenhougen, K., & Quah, S. R. (2010). *Epidemiology and demography in public health*. Academic Press.
- Kimera, E., Vindevogel, S., Reynaert, D., Justice, K. M., Rubaihayo, J., De Maeyer, J., . . . Bilsen, J. (2020). Experiences and effects of HIV-related stigma among youth living with HIV/AIDS in Western Uganda: A photovoice study. *Plos one*, 15(4), e0232359.
- Meemon, N., Paek, S. C., Yenchai, D., & Wan, T. T. (2016). Application of the WHOQOL-HIV-BREF questionnaire in HIV-infected Thai patients: reliability and validity of the instrument. *Journal of the Association of Nurses in AIDS Care*, 27(5), 698-708.
- Mrus, J. M., Leonard, A. C., Yi, M. S., Sherman, S. N., Fultz, S. L., Justice, A. C., & Tsevat, J. (2006). Health-related quality of life in veterans and nonveterans with HIV/AIDS. *Journal of general internal medicine*, 21(S5), S39-S47.
- Mutabazi-Mwesigire, D., Katamba, A., Martin, F., Seeley, J., & Wu, A. W. (2015). Factors that affect quality of life among people living with HIV attending an urban clinic in Uganda: a cohort study. *Plos one*, 10(6), e0126810.
- Ndikuno, C., Namutebi, M., Kuteesa, J., Mukunya, D., & Olwit, C. (2016). Quality of life of caregivers of patients diagnosed with severe mental illness at the national referral hospitals in Uganda. *BMC psychiatry*, 16(1), 1-9.
- Ng, L. C., Stevenson, A., Kalapurakkel, S. S., Hanlon, C., Seedat, S., Harerimana, B., . . . Koenen, K. C. (2020). National and regional prevalence of posttraumatic stress disorder in sub-Saharan Africa: A systematic review and meta-analysis. *PLoS medicine*, 17(5), e1003090.
- Organization, W. H. (1996). WHOQOL-BREF: introduction, administration, scoring and generic version of the assessment: field trial version, December 1996: World Health Organization.
- Peterman, A. H., Fitchett, G., Brady, M. J., Hernandez, L., & Cellier, D. (2002). Measuring spiritual well-being in people with cancer: the functional assessment of chronic illness therapy—Spiritual Well-being Scale (FACIT-Sp). *Annals of Behavioral Medicine*, 24(1), 49-58.
- Reid, N. G., & Boore, J. R. (1991). *Research methods and statistics in health care*: Edward Arnold.
- Sidibé, M. HIV/AIDS Branch and the Gender, Human Rights and Culture Branch of the United Nations Population Fund (UNFPA), under the coordination and direction of Lynn Collins, Technical Adviser in the HIV/AIDS Branch.

34 Religiosity and health-related quality of life in HIV positive young adults

Szaflarski, M., Ritchey, P. N., Leonard, A. C., Mrus, J. M., Peterman, A. H., Ellison, C. G., . . . Tsevat, J. (2006). Modeling the effects of spirituality/religion on patients' perceptions of living with HIV/AIDS. *Journal of general internal medicine*, 21(5), S28-S38.

UNAIDS. (2019). Factsheet - Facts on HIV and AIDS in Uganda 2021 (Based on Data ending 31st December 2020) Retrieved February, 9, 2022, from file:///C/Users/Hp-450/AppData/Local/Temp/2020%20Uganda%20HIV%20-AIDS%20fact%20sheet,%206th%20Sept%202021.pdf

Young, K., & Clark, M. (2013). Critical Synthesis Package: Duke University Religion Index (DUREL). MedEdPORTAL, 9, 9586.